

Overlap of obsessive and posttraumatic symptoms: A systematic review

José Carlos Medina Rodríguez,¹ Elia Mireya Solís Villegas¹

¹ Departamento de Enseñanza, Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz, Ciudad de México, México.

Correspondence:

José Carlos Medina Rodríguez
Departamento de Enseñanza, Instituto Nacional de Psiquiatría Ramón de la Fuente Muñiz.
Calz. México-Xochimilco 101, San Lorenzo Huipulco, Tlalpan, 14370, Ciudad de México, México.
Phone: +52 55 4443-8609.
Email: jcmedinar@imp.edu.mx

Received: 10 February 2021

Accepted: 28 June 2021

Citation:

Medina Rodríguez, J. C., & Solís Villegas, E. M. (2022). Overlap of obsessive and posttraumatic symptoms: A systematic review. *Salud Mental, 45*(3), 135-143.

DOI: [10.17711/SM.0185-3325.2022.018](https://doi.org/10.17711/SM.0185-3325.2022.018)



ABSTRACT

Background. Current medical literature suggests a symptomatic overlap between posttraumatic stress disorder (PTSD) and obsessive-compulsive disorder (OCD), which makes understanding their impact on the treatment process a challenging undertaking. **Objective.** The key aim of this work was to deliver a comprehensive overview of empirical and observational studies concerning the overlap between both psychopathologies. **Method.** Two independent authors searched PubMed, PsycINFO, and Web of Science databases between April and July 2020 utilizing relevant MeSH terms. Subsequently, a systematic scoping review was undertaken according to the PRISMA-ScR Statement to identify all relevant publications concerning the overlap of OCD and PTSD symptomatology. We included peer reviewed studies published in any language that contributed quantitative or qualitative data that addressed the relationship between OCD and PTSD symptoms as the main aim or outcome of the study. Non-peer-reviewed articles, studies that did not address this phenomenon of interest, and gray literature documents were excluded. This review was not registered into the International Prospective Register of Systematic Reviews, since, according to the National Institute of Health Research, scoping reviews are ineligible for registration into PROSPERO. **Results.** Twenty-five relevant studies were included (11 cross-sectional studies, 5 longitudinal-prospective studies, 4 case series, 4 case studies and 1 retrospective study). We did not conduct any statistical analysis due to the diversity of the included studies, thus proceeding to realize a thematic synthesis of the obtained data. **Discussion and conclusion.** A symptomatic overlap exists between both comorbidities in some clinical populations, and this relationship may hamper treatment outcomes.

Keywords: Humans, comorbidity, obsessive compulsive disorder, post-traumatic stress disorder.

RESUMEN

Antecedentes. La literatura médica sugiere la existencia de una superposición sintomática entre el trastorno por estrés posttraumático (TEPT) y el trastorno obsesivo-compulsivo (TOC), situación que puede comprometer el tratamiento de esta población clínica. **Objetivo.** El objetivo clave de este trabajo consiste en ofrecer una visión general de estudios empíricos y observacionales sobre la superposición entre ambas psicopatologías. **Método.** Dos autores independientes buscaron en las bases de datos PubMed, PsycINFO y Web of Science entre abril y julio de 2020 utilizando DeCS relevantes. Se realizó una revisión sistemática exploratoria de acuerdo con la Declaración PRISMA-ScR para identificar todas las publicaciones relevantes sobre esta superposición sintomática. Se incluyeron estudios revisados por pares publicados en cualquier idioma que aportaron datos cuantitativos o cualitativos, que abordaron la relación entre los síntomas de TOC y TEPT. Se excluyeron los artículos no revisados por pares, aquellos que no abordaron este fenómeno de interés y la literatura gris. Esta revisión no se registró en el Registro Prospectivo Internacional de Revisiones Sistemáticas, ya que acorde al Instituto Nacional de Investigación en Salud, las revisiones sistemáticas exploratorias no son elegibles para su registro en PROSPERO. **Resultados.** Se incluyeron 25 estudios relevantes (11 estudios transversales, 5 estudios longitudinales-prospectivos, 4 series de casos, 4 estudios de casos y 1 estudio retrospectivo). No se realizó ningún análisis estadístico debido a la diversidad de los estudios incluidos, por lo que se procedió a realizar una síntesis temática. **Discusión y conclusión.** Existe una superposición sintomática entre ambas comorbilidades y esta relación puede obstaculizar el tratamiento.

Palabras clave: Humanos, comorbilidad, trastorno obsesivo-compulsivo, trastorno por estrés posttraumático.

BACKGROUND

The current taxonomic framework for psychiatric disorders presents both posttraumatic stress disorder and obsessive-compulsive disorder as categories characterized by specific criteria. Nonetheless, there is a noticeable overlap in the symptomatology of these disorders, which our current diagnostic model frequently fails to recognize. Ultimately, this ends up being one of the main drawbacks associated with a categorical approach in psychiatric practice. (Kendall, Brady, & Verduin, 2001). Taking this into account, the broad spectrum of anxiety disorders is no exception to this issue. Several symptomatic domains characterized by intrusive, repetitive, and multimodal cognitions that cause distress outline this heterogeneous group of constructs. These could be further formulated into five symptomatic domains: fear, recurrent thoughts, intrusive images, physical symptoms, and avoidance (Cohen, Mychailyszyn, Settapani, Crawley, & Kendall, 2011). The obsessive-compulsive disorder (OCD) is an entity defined by the presence of obsessions and/or compulsions. Obsessions represent recurrent and persistent thoughts, impulses, or images that are intrusive, inappropriate, and experienced as distressing, while compulsions indicate repetitive, intentional behaviors carried out according to inflexible rules to reduce such distress. Similarly, the posttraumatic stress disorder (PTSD) is characterized by a cluster of symptoms that develop in response to trauma, an experience in which an individual lives or witnesses an event involving actual or threatened death, serious injury or peril to the integrity of itself or others (American Psychiatric Association, 2013). Coincidentally, OCD and PTSD share some clinical domains, with the occurrence of intrusive symptoms being common in both conditions. Whereas OCD patients tend to experience intrusive phenomena of distressing events, PTSD patients usually experience them as recurrent memories of their traumatic experience (Morina et al., 2016).

This theoretical, clinical, and empirical overlapping of functional connections between OCD and PTSD has been discussed previously (Gershuny, Baer, Radomsky, Wilson, & Jenike, 2003). However, the frequency of this phenomenon remains relatively understudied (Huppert et al., 2005; Nacasch, Fostick, & Zohar, 2011). Furthermore, similarities between recurrent, intrusive thoughts between both entities continue to be among the most challenging aspects of the clinical differential diagnosis process in current psychiatric practice (Cohen et al., 2011). Previous efforts that sought to explain these phenomena pondered a biological and environmental role following an exposure to traumatic events and OCD (Fostick, Nacasch, & Zohar, 2012).

According to this, some researchers believe that trauma in adults could be an arbiter for gene-environment transactions that lead to a late phenotypic expression of OCD (Lafleur et al., 2011; Sasson et al., 2005). Other authors suggest

that there may be a temporal connection between specific types of trauma and the development of OCD symptoms (Fontenelle et al., 2012). Likewise, interpersonal traumas and certain personality traits also may play a predictive role in PTSD diagnosis or severity in OCD patients (Gershuny et al., 2008). Interestingly, traumatic experiences may be highly prevalent in children with OCD (Lafleur et al., 2011), and child abuse may be a relevant factor in regards to clinical, functional, and trauma-related differences between pretraumatic and posttraumatic OCD diagnosis in PTSD patients (Araújo et al., 2018). In addition, mental imagery, another main feature of many anxiety disorders, has been functionally linked to earlier adverse events (Speckens, Hackmann, Ehlers, & Cuthbert, 2007). This shouldn't be surprising, given that both disorders interact with elements of threat conditioning and fear extinction, features which play a role in the development of both disorders (McGuire et al., 2016). Also, it has been shown that individuals with comorbid OCD and PTSD have a higher prevalence of severe OCD symptoms in a lifespan course than those without lifetime PTSD (Ojserkis et al., 2017). Given these points, it is worth noting that current medical literature available in regards to the relationship between PTSD and OCD is limited, which makes understanding their comorbidity and impact on their treatment outcomes a challenging undertaking (Van Kirk, Fletcher, Wanner, Hundt, & Teng, 2018). Hence, as a main objective, this research aims to provide a comprehensive overview of empirical and observational studies involving the symptomatic overlap between OCD and PTSD.

METHOD

Data source, search strategy, and research questions

No previous systematic review that focused on this phenomenon of interest was found after a comprehensive search in the PROSPERO database. Thus, this review was not registered in the International Prospective Register of Systematic Reviews, since according to the National Institute of Health Research, scoping reviews are ineligible for registration into PROSPERO. A review of literature was undertaken to identify all relevant publications concerning the overlap of OCD and PTSD symptomatology. We followed the PRISMA-ScR Statement (Moher et al., 2009), and the following research questions were formulated: 1. Does a functional connection exist between OCD and PTSD symptoms? 2. Which variables (e.g., risk factors, clinical similarities, treatment implications, etc.) influence the treatment outcomes of patients with mutual OCD and PTSD symptom clusters? Consequently, two independent authors searched PubMed, PsycINFO, and Web of Science databases between April 01 and July 26, 2020. The search was made from inception through

July 26, 2020 using the following terms: (overlap of) AND (symptoms OR symptomatology); (OCD) OR (obsessive OR compulsive) AND (PTSD) OR (posttraumatic OR trauma); (“humans”[Mesh]AND “comorbidity”[Mesh]); (“Obsessive-Compulsive Disorder / diagnosis”[Mesh]) AND “Obsessive-Compulsive Disorder / epidemiology”[Mesh] AND “Obsessive-Compulsive Disorder / psychology” [Mesh]; (“Stress Disorders, Post-Traumatic / diagnosis”[Mesh]) AND (“Stress Disorders, Post-Traumatic / epidemiology”[Mesh] AND “Stress Disorders, Post-Traumatic / psychology” [Mesh]).

Eligibility criteria

Initially, we restricted our literature search to include articles that met the following criteria according to the key words mentioned above: studies published in English in peer reviewed journals that contributed quantitative or qualitative data and that specifically addressed the relationship between OCD and PTSD symptoms as the main aim or outcome of the study. Secondly, non-peer-reviewed articles, studies that did not address this phenomenon of interest and gray literature documents were excluded. Thirdly, we eliminated all duplicate findings from these searches following a title and abstract review. Lastly, the reference lists of all articles that were relevant, as well as any recent review papers that examined this phenomenon, were surveyed in full detail to determine if they met our inclusion criteria. No other conditions than those previously described were applied to our study selection. Most initial search results (2,784) were excluded because of unrelatedness in the title or abstract regarding our subject matter (e.g., “traumatic brain injury,” “surgical trauma,” “compulsive exercise” etc.), and thus, the search resulted in 103 articles for further review. After additional inquiry, we narrowed these articles down to 25 relevant to the present study. Data and biases from these studies were scrutinized using critical appraisal tools, such as the Newcastle-Ottawa Scale for cross sectional, retrospective and cohort studies, and the Joanna Briggs Institute’s Critical Appraisal Tools for case series and case reports (Jordan, Lockwood, Munn, & Aromataris, 2019; Stang, 2010). Pertaining to their methodological approach, we included 11 cross-sectional studies, five longitudinal-prospective studies, four case series, four case studies and one retrospective study. Ultimately, all included studies were read, and key pieces of information were extracted, considering their sample size, recruitment process, design, outcomes, measures, psychometric tools used, main results, and consequences for future research.

Analysis

We did not conduct any statistical test or meta-analysis because of methodological disparities between the included

studies. Likewise, weighting the studies was not possible or done because articles with distinct methodological approaches and properties were utilized. Taking this into account, both authors chose to examine the extent, range, and characteristics of the found evidence of this phenomenon of interest. This effort meant to summarize findings from this body of knowledge that is heterogeneous in its methods and facilitated the identification of gaps in the literature to aid the planning and commissioning of future research. Thus, a thematic synthesis was done employing the data extracted, since scoping reviews is a type of knowledge synthesis that follows a systematic approach to further map evidence on a topic, helping to identify main concepts, theories, and knowledge gaps that may inspire further innovative research (Figure 1).

RESULTS

Twenty-five studies met the inclusion criteria for this review (Table 1). To our knowledge, this is the first review article addressing the putative relationship and overlap between OCD and PTSD symptom clusters utilizing studies of varied methodologies (e.g., case series, cross-sectional studies, retrospective, and prospective studies, etc.). Four types of




Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of the review process.

Table 1
Characteristics of the 25 included studies for the present review

Study authors, and year	Country	Design	Diagnosis population	Age	Gender	No. of patients	Outcome
Pitman, 1993	United States	Case study	Military personnel	46 years old	Male	1	Concurrent OCD and PTSD.
Langlois et al., 2000	Canada	Cross-sectional study	Students	22.5 years old (mean age)	Male and female	254	Disfunction caused by worry is more severe than disfunction caused by obsessions.
de Silva & Marks, 2001	United States	Case series	Treatment resistant OCD patients ^a	16-76 years old	Male and female	104	82% reported a history of at least one trauma, 49.4% met criteria for PTSD (criteria non-disclosed).
Gershuny et al., 2002	United States	Cross-sectional study	OCD patients (DSM-IV ^b) with previous failed trials of behavior therapy and/or medication	19-66 years old	Male and female	15	Patients with comorbid PTSD had worse outcomes in behavior therapy.
Gershuny et al., 2003	United States	Case series	Patients who met criteria for OCD (YBOCS ^c) and PTSD based on interview and self-report	Not reported	Male and female	4	An inverse relationship between PTSD symptoms versus OCD symptoms that may hamper therapy effectiveness.
Huppert et al., 2005	United States	Cross-sectional study	128 OCD patients (OCI-R ^d), 109 PTSD patients (PDS ^e), 63 patients with other anxiety disorders and 40 college students	21.7-44.5 years old (mean age)	Male and female	340	Overlap of OCD and PTSD symptoms using continuous rating scales.
Sasson et al., 2005	Israel	Case series	Military personnel	Not reported	Male	13	4 Case reports had evidence of previous trauma regarding OCD symptom onset.
Cromer et al., 2007	United States	Cross-sectional	OCD patients (YBOCS ^c)	41 years old (mean age)	Male and female	265	Traumatic life events are associated with higher OCD severity in Y-BOCS ^c severity ratings.
Grabe et al., 2007	Germany	Cross-sectional study	OCD patients (DSM-IV ^b)	37.4 years old (mean age)	Male and female	210	No apparent relationship between trauma, PTSD and OCD.
Speckens et al., 2007	United Kingdom	Prospective study	OCD patients (DSM-IV ^b)	35.7 years old (mean age)	Male and female	37	Patients with traumatic mental images had more OCD symptoms.
de Moraes et al., 2008	Brazil	Case study	PTSD patients (criteria non-disclosed)	35 years old	Male	1	Scored 30 on the initial Y-BOCS ^c symptom severity post-trauma and non-responsive to pharmacotherapy.
Ehring et al., 2008	United Kingdom	Prospective study	Motor Vehicle Accident Survivors	35.17 years old (mean age)	Male and female	147	22.4 % participants met criteria for PTSD (DSM-IV ^b) and alongside with checking behaviors.
Gershuny et al., 2008	United States	Cross-sectional study	OCD patients (DSM-IV-TR)	32 years old (mean age)	Male and female	104	82% reported at least one trauma, 39.4% met criteria for PTSD.
Shavitt et al., 2010	Brazil	Prospective study	OCD patients (DSM-IV ^b)	36.3 years old (mean age)	Male and female	219	Patients with OCD and PTSD presented a greater response in symptom dimensions through therapy versus patients without PTSD.
Nacasch et al., 2011	Israel	Cross-sectional study	PTSD patients (MINI ^f)	44.29 years old (mean age)	Male and female	44	41% of PTSD patients had comorbid OCD (MINI ^f).
Badour et al., 2012	United States	Cross-sectional study	Women with diagnosis of previous sexual trauma or physical assault (DSM-IV-TR)	28.37 years old (mean age)	Female	49	Peritraumatic self-focused disgust was significantly related to contamination-based OCD symptoms (OCI-R ^d).

Table 1 (continued)

Study authors, and year	Country	Design	Diagnosis population	Age	Gender	No. of patients	Outcome
Fontenelle et al., 2012	Brazil	Retrospective study	OCD patients (DSM-IV ^a)	37.3 years old (mean age)	Male and female	1001	OCD occurring after or near the onset of PTSD is associated with distinct clinical features specially when patients did not had history of pre-traumatic OCD.
Fostick et al., 2012	Israel	Case series	Military personnel	21 years old	Male	5	The onset of PTSD and OCD was simultaneous and occurred after combat trauma.
Semiz et al., 2014	Turkey	Cross-sectional study	OCD patients (DSM-IV ^a)	35.80 years old (mean age)	Female	120	Dissociation (according to DES ^b) scores may be a predictor of poor treatment outcome.
Morina et al., 2016	Switzerland	Cross-sectional study	Civilian survivors of Kosovo War.	43.0 years old (mean age)	Male	51	Participants with high levels of PTSD symptoms (PDS ^c) were significantly more likely to have OCD symptoms (OCI-R ^d).
Ojserkis et al., 2017	United States	Prospective study	OCD patients (DSM-IV ^a)	39.16 years old (mean age)	Male and female	266	PTSD patients with comorbid OCD reported significantly more OCD symptoms (Y-BOCS ^e).
Araujo et al., 2018	Brazil	Prospective study	Patients with comorbid PTSD and OCD (DSM-IV ^a)	40.1 years old (mean age)	Male and female	63	45.2% of PTSD patients with pre-traumatic OCD had history of childhood abuse.
Van Kirk et al., 2018	United States	Case study	Military personnel	Mid-40s	Male	1	Apparent functional overlap between OCD (Y-BOCS ^e) and PTSD (PCL-5 ^f) symptoms.
Franklin & Raines, 2019	United States	Cross-sectional study	Military personnel	45.1 years old (mean age)	Male and female	117	Significant overlap between PTSD and OCD symptoms according to self-report instruments (PCL-5 ^f , DOCS ^g).
Rossi et al., 2020	Italy	Case study	Motor Vehicle Accident survivor	35 years old	Male	1	OCD diagnosis (DSM-5 ^h) with apparent treatment response to sertraline – aripiprazole.

Notes:

- ^a Treatment resistant OCD patient: Participants with failure of at least one adequate prior treatment trial or at least two types of medication trials.
- ^b DSM-IV: Diagnostic and Statistical Manual of Mental Disorders Fourth Edition.
- ^c YBOCS: Yale Brown Obsessive Compulsive Scale.
- ^d OCI-R: Obsessive Compulsive Inventory-Revised.
- ^e PDS: Posttraumatic Diagnostic Scale.
- ^f DSM-IV-TR: Diagnostic and Statistical Manual of Mental Disorders Fourth Edition Text Revision.
- ^g MINI: Mini International Neuropsychiatry Interview.
- ^h Dissociative Experiences Scale.
- ⁱ PCL-5: Posttraumatic Stress Disorder for DSM-5.
- ^j DOCS: Dimensional Obsessive-Compulsive Scale.
- ^k DSM-5: Diagnostic and Statistical Manual of Mental Disorders Fifth Edition.

research were identified through the thematic analysis of the included studies: clinical domains; risk factors and other correlates; prognostic factors; and treatment approaches. Thus, our results section describes each of these categories. To establish confidence in the evidence reviewed we utilized the ConQual approach, a practical tool developed by the Joanna Briggs Institute, to grade the included articles according to the type of study, dependability, and credibility. The proposed system would then give an overall score of High, Moderate, Low, or Very Low. This ranking can be considered a rating of confidence in the synthesized findings. Afterwards, a summary of findings table was developed to convey the key findings of the review in a tabular format with the aim of improving the accessibility and usefulness of the systematic scoping review (Table 2; Jordan et al., 2019; Stang, 2010).

Clinical domains ($n = 7$)

Several of the included studies addressed phenomenological and symptomatic domains of OCD and PTSD symptoms in a variety of community samples. These reports were mostly comprised of observational studies or clinical cases. To start with, relatively recent literature has showcased PTSD patients that manifest cognitions such as compulsions that have been described to occur strictly within the context of trauma, and that these may not be necessarily indicative of a comorbid diagnosis of OCD (de Silva & Marks, 2001). Similarly, other researchers have noticed that adults with in-

trusive cognitions tend to be more aware of specific external or internal stimuli in comparison to other adults with obsessions. Also, it was observed that *worry*, as a dimensional cognition, tends to be more severe than dysfunction-generating *obsessions* (Langlois, Freeston, & Ladouceur, 2000). Coincidentally, *mental images*, a recurrent phenomenon in many anxiety disorders, tend to be portrayed as irrational in nature in OCD patients. In comparison, it is important to note that individuals with this diagnosis can also experience intrusive images associated with earlier trauma (Speckens et al., 2007). Likewise, *checking behaviors*, another symptomatic dimension in obsessive and anxiety disorders, have been noted in a sample of motor vehicle accident survivors with up to six months post-trauma (Ehring, Ehlers, & Glucksman, 2008). In regards to a possible temporal causality factor, another study found that nearly all OCD symptoms were more frequent or severe among posttraumatic OCD patients, with the exceptions of certain symptomatic domains, such as symmetry-seeking behaviors, hand washing, and hypersensitivity to specific sensory phenomena, and that these symptoms were more common in pretraumatic OCD (Fontenelle et al., 2012). In the same way, another study about civilian survivors of war found that compulsive washing was the most frequent and severe symptom in participants with high levels of PTSD symptoms (Morina et al., 2016). Likewise, some researchers found a similar thematic connection in an outpatient sample of veterans with OCD self-report. These results may show that some OCD symptom profiles could be similar to PTSD cognitive symptom

Table 2
Confidence of the synthesized findings graded according to the Joanna Briggs Institute's ConQual approach

Population: Participants with overlapping clinical symptoms of both OCD and PTSD.

Phenomena of interest: Overlap of OCD and PTSD symptomatology.

Context: Participants of any age, sex, ethnicity, and culture that manifest comorbid OCD and PTSD symptoms

<i>Synthesized finding</i>	<i>Type of research</i>	<i>Dependability</i>	<i>Credibility</i>	<i>ConQual score</i>
<i>Clinical domains</i> There seems to be a temporal and causal factor relating to the specific context and timing of trauma, since some symptom clusters manifest differently according to the moment of experienced trauma.	Observational and empirical	Downgraded one level	Downgraded one level	Moderate
<i>Risk factors and other correlates</i> Of interest, early life adversity such as sexual and interpersonal violence and other stressors like combat trauma seem to influence the development of both OCD and PTSD symptoms.	Observational and empirical	Downgraded one level	Downgraded one level	Moderate
<i>Prognostic factors</i> It appears that when both psychopathologies concur that the overall prognosis is diminished.	Observational and empirical	Downgraded one level	Downgraded one level	Moderate
<i>Treatment approaches</i> To date, no single clinical guideline or evidence-based recommendation exists that focuses on both entities when comorbid. Some reports show mixed results when employing pharmacological and psychosocial approaches	Observational and empirical	Downgraded two levels	Downgraded two levels	Low

Note: The downgrade of the main scores was due to common dependability issues across the included studies and owing to a mix of equivocal and unequivocal findings. For example, in most studies, the researchers had no acknowledgement of their influence on their or future research.

clusters (e.g., feeling responsible for harm, and unacceptable violent, or sexual thoughts; Franklin & Raines, 2019).

Risk factors and other correlates ($n = 9$)

Several findings included in this review discussed the relevance of specific stressors that resulted in the development of OCD symptoms in PTSD patients. Of interest, an old case report describes an individual without previous overt psychopathology who, after experiencing combat stressors, underwent development of concurrent OCD and PTSD, showing classical symptoms, such as obsessions of violent nature, contamination, doubt, compulsive hand-washing, cleaning, checking, counting, touching, attempts to resist these compulsions, and other related phenomena (Pitman, 1993). Subsequently, another series of cases suggested that the course of each disorder may run a heterotypical course when comorbid, and that this may address a different, heterogeneous subtype of OCD resulting from trauma (Sasson et al., 2005). Also, in one of the largest studies to date, 52% of 365 individuals with previous OCD had at least one or more earlier traumatic events (Cromer, Schmidt, & Murphy, 2007). In contrast, a study found no apparent relationship between trauma, acute stress disorder, PTSD, or symptoms in patients with OCD when compared to control patients from the general population (Grabe et al., 2007). Afterwards, a brief report observed that a high proportion of individuals seeking treatment for resistant OCD showcased earlier history of childhood and adulthood interpersonal violence. Of these, incestuous childhood sexual abuse, witnessing violence, and a greater frequency of experienced traumatic experiences were taken into account as putatively predictive of higher PTSD severity (Gershuny et al., 2008). Ratifying the findings of this report, another study found that a history of childhood abuse was statistically more common amidst PTSD patients with pre-traumatic OCD than among their counterparts with post-traumatic OCD (Araújo et al., 2018). Other researchers found in their study sample that nearly half of their participants developed OCD symptoms following exposure to combat- or terror- related trauma, and that most of those eventually met criteria for full OCD (Nacasch et al., 2011). In the same vein, a case series of five Israeli veterans with no prior personal or family history of OCD or other psychiatric comorbidity who were diagnosed simultaneously with PTSD and OCD following combat trauma, described that OCD symptoms were initially related to the previous trauma, but then expanded to include other non-traumatic related triggers (Fostick et al., 2012). Incidentally, another group of investigators that studied peritraumatic fear, self-focused and other-focused disgust found out that self-focused disgust was related to contamination-based OCD symptoms, while peritraumatic fear and other-focused disgust was not significantly related with this type of phenomena (Badour, Bown, Adams, Bunaciu, & Feldner, 2012).

Prognostic factors ($n = 6$)

A small case series noted an apparent inverse relationship between symptom intensity in both conditions (when OCD symptoms lessen, symptoms of PTSD increase, and vice versa.). Likewise, the authors of this work discussed that targeting OCD (and perhaps PTSD) in isolation may hamper therapy effectiveness (Gershuny et al., 2003). Moreover, another study found a significant relationship between OCD and PTSD symptoms in clinical and non-clinical samples utilizing continuous ratings instruments like the Posttraumatic Diagnostic Scale and the Obsessive-Compulsive Inventory (Huppert et al., 2005). Of interest, another clinical case described a patient without earlier psychiatric comorbidities who developed PTSD, depression, and OCD after a work-related trauma. This patient scored 30 on the initial Yale Brown Obsessive-Compulsive Scale (YBOCS) and 24 on the Beck Depression Inventory. Then, he reported panic attack phenomena in trauma-related situations or when his OCD symptoms worsened. Curiously, the authors of this report commented that adequately dosed medications provided no response, including the use of fluoxetine, sertraline, citalopram, bupropion, paroxetine, and fluvoxamine (de Moraes, Torresan, Trench, & Torres, 2008). Another study that compared the clinical characteristics of OCD with and without comorbid PTSD in an outpatient sample also revealed a high prevalence of OCD symptom severity for the comorbid group when the severity was measured using multiple OCD symptom dimensions (Shavitt et al., 2010). Other researchers noticed that severe OCD symptoms were found in patients with a higher degree of dissociative symptoms. In this report, it was shown that YBOCS scores were significantly correlated to severity of dissociation, anxiety, depression, and traumatic phenomena (Semiz, Inanc, & Bezgin, 2014). Lastly, a naturalistic, seven-year, longitudinal OCD study reported that individuals with comorbid OCD and PTSD had more severe and impairing OCD symptoms, poorer insight, lower quality of life, and greater incidences of lifetime affective and substance use disorders at the baseline assessment (Ojserkis et al., 2017).

Treatment approaches ($n = 3$)

Although scarce, this section addresses few treatment modalities employed, or pondered, in patients with both comorbidities. Even if unclear, some studies have found that behavioral treatment of OCD (with or without pharmacological interventions) may be hampered by the presence of comorbid PTSD and may even be contraindicated, since this type of interventions have been shown to reactivate symptoms associated with the traumatic event (Gershuny, Baer, Jenike, Minichiello, & Wilhelm, 2002). Some researchers have also distinguished in a case series that an integrated treatment approach to address both OCD and PTSD symp-

toms through a cognitive-behavioral system (cognitive restructuring to modify trauma-related beliefs paired with exposure and prevention therapy for OCD symptoms) may be effective when there is a functional connection between both entities (Van Kirk et al., 2018). Lastly, a recent case study reported an apparent successful treatment approach with aripiprazole augmentation of sertraline (Rossi, Niolu, Siracusano, Rossi, & Di Lorenzo, 2020).

DISCUSSION AND CONCLUSION

The aim of this work was to provide a comprehensive overview of empirical and observational research concerning the overlap between OCD and PTSD symptoms. The studies reviewed indicate that a functional relationship between trauma and the development of both PTSD and OCD symptoms in some clinical populations may exist (Knowles, Sripada, Defever, & Rauch, 2019). It should be noted that some reports have stated that the clinical features of OCD symptoms vary according to the timing of the traumatic experience, suggesting the possibility of a subtype of OCD (pretraumatic vs posttraumatic; Araújo et al., 2018). Also, in phenomenological terms, many studies have observed that some symptom clusters correlate with the natural history and severity of both comorbidities. For example, cognitions such as worry, which are common in both disorders, may be predictive of higher disfunction when compared to isolated OCD symptoms (Langlois et al., 2000). Likewise, dissociation, a common symptom in trauma survivors, may be a predictor of a poorer treatment outcome in patients with OCD (Semiz et al., 2014). Peritraumatic self-disgust was significantly related to contamination cognitions, and war survivors have been shown to have high prevalence of OCD symptoms (Badour et al., 2012). This functional overlap has been consistently observed in multiple reports and measured through validated and standardized self-report instruments. These findings have utmost importance regarding treatment outcomes, since this has been exemplified in a few studies that highlight the difficulties of managing both disorders when comorbid.

Thus, as a conclusion, we propose a functional connection between the symptoms of both disorders. This connection may be of considerable difficulty to distinguish when comorbid and could hamper the prognosis of some patients. Hence, we recommend that future research should cover this phenomenon in detail. These efforts may aim to optimize treatment modalities and focus on specific outcomes for this patient population that carries a great burden and, perhaps, may be underdiagnosed. More longitudinal and qualitative methodologies could be combined to offer a comprehensive picture of this phenomenon, taking into consideration the sociodemographic and psychosocial environment it is part of.

We would like to add that this revision was expanded from previous research and targeted specific outcomes of studies covering this topic, but it is not without its limitations. Firstly, data regarding this phenomenon is scarce, and the few reports that were included may be highly heterogeneous in their demographical population and methodological approaches. Distinct instruments or measures to diagnose both disorders and their symptoms were employed in many studies, which limited our capacity to aggregate results. There was also an imbalance in sample sizes in the included reports, and therefore, we were limited to make any statistical comparisons across study groups. Secondly, while this phenomenon may be an emerging area of interest, it should be noted that the assumptions derived from this work cannot explain the complex mechanisms in which both disorders connect and manifest, which should also be a focus of future research. Thirdly, while we followed rigorous search methods to identify relevant papers, the review was limited only to published reports that we were able to locate. Lastly, the exclusion of research in other languages should also be taken in consideration even though our search aimed to include studies in any vernacular, making it possible that additional data that covers this topic exists but was not included in this review (i.e., because of publication bias). As a concluding remark, we would like to issue the possible overlap of symptoms in these disorders, seeing that both are associated with great burden and may be underdiagnosed.

Funding

None.

Conflict of interest

The authors declare they have no conflicts of interest.

Acknowledgements

We express great appreciation to Dr. Ilyamín Merlín García for his valuable supervision and constructive suggestions during the planning and development of this review.

REFERENCES

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th edition). Washington, DC: American Psychiatric Publishing.
- Araújo, A. X. G. de, Fontenelle, L. F., Berger, W., Luz, M. P. da, Pagotto, L. F. A. da C., Marques-Portella, C., ... Mendlowicz, M. V. (2018). Pre-traumatic vs post-traumatic OCD in PTSD patients: Are differences in comorbidity rates and functional health status related to childhood abuse? *Comprehensive Psychiatry*, 87, 25-31. doi: 10.1016/j.comppsych.2018.08.012
- Badour, C. L., Bown, S., Adams, T. G., Bunaciu, L., & Feldner, M. T. (2012). Specificity of fear and disgust experienced during traumatic interpersonal victimization in predicting posttraumatic stress and contamination-based obsessive-compulsive symptoms. *Journal of Anxiety Disorders*, 26(5), 590-598. doi: 10.1016/j.janxdis.2012.03.001
- Cohen, J., Mychailyszyn, M., Settiani, C., Crawley, S., & Kendall, P. C. (2011). Issues in Differential Diagnosis: Considering Generalized Anxiety Disorder, Obsessive-Compulsive Disorder, and Posttraumatic Stress Disorder. In

- Handbook of Child and Adolescent Anxiety Disorders* (pp. 23-35). New York: Springer. doi: 10.1007/978-1-4419-7784-7_3
- Cromer, K. R., Schmidt, N. B., & Murphy, D. L. (2007). An investigation of traumatic life events and obsessive-compulsive disorder. *Behaviour Research and Therapy, 45*(7), 1683-1691. doi: 10.1016/j.brat.2006.08.018
- de Moraes, E. C., Torresan, R. C., Trench, E. V., & Torres, A. R. (2008). A possible case of posttraumatic obsessive-compulsive disorder. *Revista Brasileira de Psiquiatria, 30*, 291. doi: 10.1590/S1516-44462008000300018
- de Silva, P., & Marks, M. (2001). Traumatic experiences, post-traumatic stress disorder and obsessive-compulsive disorder. *International Review of Psychiatry, 13*(3), 172-180. doi: 10.1080/09540260120074037
- Ehring, T., Ehlers, A., & Glucksman, E. (2008). Do Cognitive Models Help in Predicting the Severity of Posttraumatic Stress Disorder, Phobia, and Depression After Motor Vehicle Accidents? A Prospective Longitudinal Study. *Journal of Consulting and Clinical Psychology, 76*(2), 219-230. doi: 10.1037/0022-006X.76.2.219
- Fontenelle, L. F., Cocchi, L., Harrison, B. J., Shavitt, R. G., do Rosário, M. C., Ferrão, Y. A., ... Torres, A. R. (2012). Towards a post-traumatic subtype of obsessive-compulsive disorder. *Journal of Anxiety Disorders, 26*(2), 377-383. doi: 10.1016/j.janxdis.2011.12.001
- Fostick, L., Nacasch, N., & Zohar, J. (2012). Acute obsessive compulsive disorder (OCD) in veterans with posttraumatic stress disorder (PTSD). *World Journal of Biological Psychiatry, 13*(4), 312-315. doi: 10.3109/15622975.2011.607848
- Franklin, C. L., & Raines, A. M. (2019). The overlap between OCD and PTSD: Examining self-reported symptom differentiation. *Psychiatry Research, 280*, 112508. doi: 10.1016/j.psychres.2019.112508
- Gershuny, B. S., Baer, L., Jenike, M. A., Minichiello, W. E., & Wilhelm, S. (2002). Comorbid posttraumatic stress disorder: Impact on treatment outcome for obsessive-compulsive disorder. *American Journal of Psychiatry, 159*(5), 852-854. doi: 10.1176/appi.ajp.159.5.852
- Gershuny, B. S., Baer, L., Parker, H., Gentes, E. L., Infield, A. L., & Jenike, M. A. (2008). Trauma and posttraumatic stress disorder in treatment-resistant obsessive-compulsive disorder. *Depression and Anxiety, 25*(1), 69-71. doi: 10.1002/da.20284
- Gershuny, B. S., Baer, L., Radosky, A. S., Wilson, K. A., & Jenike, M. A. (2003). Connections among symptoms of obsessive-compulsive disorder and posttraumatic stress disorder: a case series. *Behaviour Research and Therapy, 41*(9), 1029-1041. doi: 10.1016/S0005-7967(02)00178-X
- Grabe, H. J., Ruhrmann, S., Spitzer, C., Josepeit, J., Ettelt, S., Buhtz, F., ... Freyberger, H. J. (2007). Obsessive-compulsive disorder and posttraumatic stress disorder. *Psychopathology, 41*(2), 129-134. doi: 10.1159/000112029
- Huppert, J. D., Moser, J. S., Gershuny, B. S., Riggs, D. S., Spokas, M., Filip, J., ... Foa, E. B. (2005). The relationship between obsessive-compulsive and posttraumatic stress symptoms in clinical and non-clinical samples. *Journal of Anxiety Disorders, 19*(1), 127-136. doi: 10.1016/j.janxdis.2004.01.001
- Jordan, Z., Lockwood, C., Munn, Z., & Aromataris, E. (2019). The updated Joanna Briggs Institute Model of Evidence-Based Healthcare. *International Journal of Evidence-Based Healthcare, 17*(1), 58-71. doi: 10.1097/XEB.0000000000000155
- Kendall, P. C., Brady, E. U., & Verduin, T. L. (2001). Comorbidity in childhood anxiety disorders and treatment outcome. *Journal of the American Academy of Child & Adolescent Psychiatry, 40*(7), 787-794. doi: 10.1097/00004583-200107000-00013
- Knowles, K. A., Sripada, R. K., Defever, M., & Rauch, S. A. M. (2019). Comorbid mood and anxiety disorders and severity of posttraumatic stress disorder symptoms in treatment-seeking veterans. *Psychological Trauma: Theory, Research, Practice, and Policy, 11*(4), 451-458. doi: 10.1037/tra0000383
- Lafleur, D. L., Petty, C., Mancuso, E., McCarthy, K., Biederman, J., Faro, A., ... Geller, D. A. (2011). Traumatic events and obsessive compulsive disorder in children and adolescents: Is there a link? *Journal of Anxiety Disorders, 25*(4), 513-519. doi: 10.1016/j.janxdis.2010.12.005
- Langlois, F., Freeston, M. H., & Ladouceur, R. (2000). Differences and similarities between obsessive intrusive thoughts and worry in a non-clinical population: Study 1. *Behaviour Research and Therapy, 38*(2), 157-173. doi: 10.1016/S0005-7967(99)00027-3
- McGuire, J. F., Orr, S. P., Essoe, J. K.-Y., McCracken, J. T., Storch, E. A., & Piacentini, J. (2016). Extinction learning in childhood anxiety disorders, obsessive compulsive disorder and post-traumatic stress disorder: implications for treatment. *Expert Review of Neurotherapeutics, 16*(10), 1155-1174. doi: 10.1080/14737175.2016.1199276
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine, 6*(7), e1000097. doi: 10.1371/journal.pmed.1000097
- Morina, N., Sulaj, V., Schnyder, U., Klaghofer, R., Müller, J., Martin-Sölch, C., & Rufer, M. (2016). Obsessive-compulsive and posttraumatic stress symptoms among civilian survivors of war. *BMC Psychiatry, 16*(1), 115. doi: 10.1186/s12888-016-0822-9
- Nacasch, N., Fostick, L., & Zohar, J. (2011). High prevalence of obsessive-compulsive disorder among posttraumatic stress disorder patients. *European Neuropsychopharmacology, 21*(12), 876-879. doi: 10.1016/j.euroneuro.2011.03.007
- Ojserkis, R., Boisseau, C. L., Reddy, M. K., Mancebo, M. C., Eisen, J. L., & Rasmussen, S. A. (2017). The impact of lifetime PTSD on the seven-year course and clinical characteristics of OCD. *Psychiatry Research, 258*, 78-82. doi: 10.1016/j.psychres.2017.09.042
- Pitman, R. K. (1993). Posttraumatic obsessive-compulsive disorder: A case study. *Comprehensive Psychiatry, 34*(2), 102-107. doi: 10.1016/0010-440X(93)90054-8
- Rossi, R., Niolu, C., Siracusano, A., Rossi, A., & Di Lorenzo, G. (2020). A Case of Comorbid PTSD and Posttraumatic OCD Treated with Sertraline-Aripiprazole Augmentation. *Case Reports in Psychiatry, 2020*, 2616492. doi: 10.1155/2020/2616492
- Sasson, Y., Dekel, S., Nacasch, N., Chopra, M., Zinger, Y., Amital, D., & Zohar, J. (2005). Posttraumatic obsessive-compulsive disorder: A case series. *Psychiatry Research, 135*(2), 145-152. doi: 10.1016/j.psychres.2004.05.026
- Semiz, U. B., Inanc, L., & Bezgin, C. H. (2014). Are trauma and dissociation related to treatment resistance in patients with obsessive-compulsive disorder? *Social Psychiatry and Psychiatric Epidemiology, 49*(8), 1287-1296. doi: 10.1007/s00127-013-0787-7
- Shavitt, R. G., Valério, C., Fossaluza, V., da Silva, E. M., Cordeiro, Q., Diniz, J. B., ... Miguel, E. C. (2010). The impact of trauma and post-traumatic stress disorder on the treatment response of patients with obsessive-compulsive disorder. *European Archives of Psychiatry and Clinical Neuroscience, 260*(2), 91-99. doi: 10.1007/s00406-009-0015-3
- Speckens, A. E. M., Hackmann, A., Ehlers, A., & Cuthbert, B. (2007). Imagery special issue: Intrusive images and memories of earlier adverse events in patients with obsessive compulsive disorder. *Journal of Behavior Therapy and Experimental Psychiatry, 38*(4), 411-422. doi: 10.1016/j.jbtep.2007.09.004
- Stang, A. (2010). Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *European Journal of Epidemiology, 25*(9), 603-605. doi: 10.1007/s10654-010-9491-z
- Van Kirk, N., Fletcher, T. L., Wanner, J. L., Hundt, N., & Teng, E. J. (2018). Implications of comorbid OCD on PTSD treatment: A case study. *Bulletin of the Menninger Clinic, 82*(4), 344-359. doi: 10.1521/bumc.2018.82.4.344